

Topological properties of Berry's phase

Kazuo Fujikawa

*Institute of Quantum Science, College of Science and Technology
Nihon University, Chiyoda-ku, Tokyo 101-8308, Japan*

Abstract

By using a second quantized formulation of level crossing, which does not assume adiabatic approximation, a convenient formula for geometric terms including off-diagonal terms is derived. The analysis of geometric phases is reduced to a simple diagonalization of the Hamiltonian in the present formulation. If one diagonalizes the geometric terms in the infinitesimal neighborhood of level crossing, the geometric phases become trivial for any finite time interval T . The topological interpretation of Berry's phase such as the topological proof of phase-change rule thus fails in the practical Born-Oppenheimer approximation, where a large but finite ratio of two time scales is involved.

The geometric phases are mostly defined in the framework of adiabatic approximation [1]-[6], though a non-adiabatic treatment has been considered in, for example, [7] and the (non-adiabatic) correction to the geometric phases has been analyzed [8]. One may then wonder if some of the characteristic properties generally attributed to the geometric phases are the artifacts of the approximation. We here show that the topological properties of the geometric phases associated with level crossing are the artifacts of the adiabatic approximation which assumes the infinite time interval $T \rightarrow \infty$ [2]. To substantiate this statement, we start with the exact definition of geometric terms associated with level crossing. The level crossing problem is neatly formulated by using the second quantization technique without assuming adiabatic approximation. The analysis of phase factors in this formulation is reduced to a diagonalization of the Hamiltonian, and thus it is formulated both in the path integral and in the operator formulation. We thus derive a convenient formula for geometric terms [1] and their off-diagonal counter parts. (As for off-diagonal geometric phases, see [9] where the off-diagonal geometric phases in the framework of an adiabatic picture in the first quantization have been proposed, and their properties have been analyzed in [10, 11, 12].) Our formula allows us to analyze the topological properties of the geometric terms precisely in the infinitesimal neighborhood of level crossing. At the level crossing point, the conventional energy eigenvalues become degenerate but the degeneracy is lifted if one diagonalizes the geometric terms. It is then shown that the geometric phases become trivial (and thus no monopole singularity) in the infinitesimal neighborhood of level crossing for any finite time interval T . This is proved independently of the adiabatic approximation. The topological interpretation [3, 1] of geometric phases such as the topological proof of Longuet-Higgins' phase-change rule [4] thus

fails for any finite T such as in the practical Born-Oppenheimer approximation, where a large but finite ratio of two time scales is involved. In practical physical applications of geometric phases, finite T is always relevant and our analysis implies that the widely used terminology of “topological phases” for the geometric phases should be taken with great care.

We start with the generic (hermitian) Hamiltonian

$$\hat{H} = \hat{H}(\hat{\vec{p}}, \hat{\vec{x}}, X(t)) \quad (1)$$

for a single particle theory in a slowly varying background variable $X(t) = (X_1(t), X_2(t), \dots)$. The path integral for this theory for the time interval $0 \leq t \leq T$ in the second quantized formulation is given by

$$Z = \int \mathcal{D}\psi^* \mathcal{D}\psi \exp\left\{\frac{i}{\hbar} \int_0^T dt d^3x [\psi^*(t, \vec{x}) i\hbar \frac{\partial}{\partial t} \psi(t, \vec{x}) - \psi^*(t, \vec{x}) \hat{H}(\hat{\vec{p}}, \hat{\vec{x}}, X(t)) \psi(t, \vec{x})]\right\}. \quad (2)$$

We then define a complete set of eigenfunctions

$$\begin{aligned} \hat{H}(\hat{\vec{p}}, \hat{\vec{x}}, X(0)) u_n(\vec{x}, X(0)) &= \lambda_n u_n(\vec{x}, X(0)), \\ \int d^3x u_n^*(\vec{x}, X(0)) u_m(\vec{x}, X(0)) &= \delta_{nm}, \end{aligned} \quad (3)$$

and expand $\psi(t, \vec{x}) = \sum_n a_n(t) u_n(\vec{x}, X(0))$. We then have $\mathcal{D}\psi^* \mathcal{D}\psi = \prod_n \mathcal{D}a_n^* \mathcal{D}a_n$ and the path integral is written as

$$Z = \int \prod_n \mathcal{D}a_n^* \mathcal{D}a_n \exp\left\{\frac{i}{\hbar} \int_0^T dt \left[\sum_n a_n^*(t) i\hbar \frac{\partial}{\partial t} a_n(t) - \sum_{n,m} a_n^*(t) E_{nm}(X(t)) a_m(t) \right]\right\} \quad (4)$$

where

$$E_{nm}(X(t)) = \int d^3x u_n^*(\vec{x}, X(0)) \hat{H}(\hat{\vec{p}}, \hat{\vec{x}}, X(t)) u_m(\vec{x}, X(0)). \quad (5)$$

We next perform a unitary transformation $a_n = U(X(t))_{nm} b_m$ where

$$U(X(t))_{nm} = \int d^3x u_n^*(\vec{x}, X(0)) v_m(\vec{x}, X(t)) \quad (6)$$

with the instantaneous eigenfunctions of the Hamiltonian

$$\begin{aligned} \hat{H}(\hat{\vec{p}}, \hat{\vec{x}}, X(t)) v_n(\vec{x}, X(t)) &= \mathcal{E}_n(X(t)) v_n(\vec{x}, X(t)), \\ \int d^3x v_n^*(\vec{x}, X(t)) v_m(\vec{x}, X(t)) &= \delta_{n,m}. \end{aligned} \quad (7)$$

We emphasize that $U(X(t))$ is a unit matrix both at $t = 0$ and $t = T$ if $X(T) = X(0)$, and thus $\{a_n\} = \{b_n\}$ both at $t = 0$ and $t = T$. We take the time T as the period of the

slow variable $X(t)$. We can thus re-write the path integral as

$$Z = \int \prod_n \mathcal{D}b_n^* \mathcal{D}b_n \exp \left\{ \frac{i}{\hbar} \int_0^T dt \left[\sum_n b_n^*(t) i\hbar \frac{\partial}{\partial t} b_n(t) + \sum_{n,m} b_n^*(t) \langle n | i\hbar \frac{\partial}{\partial t} | m \rangle b_m(t) - \sum_n b_n^*(t) \mathcal{E}_n(X(t)) b_n(t) \right] \right\} \quad (8)$$

where the second term in the action stands for the term commonly referred to as Berry's phase[1](in the interpretation of the phase as dynamical [8]) and its off-diagonal counter part. The second term is defined by

$$\begin{aligned} (U(t)^\dagger i\hbar \frac{\partial}{\partial t} U(t))_{nm} &= \int d^3x v_n^*(\vec{x}, X(t)) i\hbar \frac{\partial}{\partial t} v_m(\vec{x}, X(t)) \\ &\equiv \langle n | i\hbar \frac{\partial}{\partial t} | m \rangle. \end{aligned} \quad (9)$$

In the operator formulation of the second quantized theory, we thus obtain the effective Hamiltonian (depending on Bose or Fermi statistics)

$$\hat{H}_{eff}(t) = \sum_n b_n^\dagger(t) \mathcal{E}_n(X(t)) b_n(t) - \sum_{n,m} b_n^\dagger(t) \langle n | i\hbar \frac{\partial}{\partial t} | m \rangle b_m(t) \quad (10)$$

with $[b_n(t), b_m^\dagger(t)]_\mp = \delta_{n,m}$. Note that these formulas (4), (8) and (10) are exact. The off-diagonal geometric terms in (10), which are crucial in the analysis below, are missing in the usual adiabatic approximation in the first quantization¹. In our picture, all the phase factors are included in the Hamiltonian, and for this reason, the terminology “geometric terms” is used for the terms in the Hamiltonian and the “geometric phases” is reserved for the geometric terms when they are explicitly interpreted as the phase factors of a specific state vector.

We are mainly interested in the topological properties in the infinitesimal neighborhood of level crossing. We thus assume that the level crossing takes place only between the

¹It is possible to show that

$$\begin{aligned} &\langle n | T^* \exp \left\{ -(i/\hbar) \int_0^T dt \hat{\mathcal{H}}_{eff}(t) \right\} | n \rangle \\ &= \langle n(T) | T^* \exp \left\{ -(i/\hbar) \int_0^T dt \hat{H}(\hat{\vec{p}}, \hat{\vec{x}}, X(t)) \right\} | n(0) \rangle \end{aligned}$$

where T^* stands for the time ordering operation. The state $|n\rangle$ on the left-hand side is defined by $b_n^\dagger(0)|0\rangle$ whereas $|n(0)\rangle$ and $|n(T)\rangle$ on the right-hand side are defined by the eigenfunctions of $\hat{H}(\hat{\vec{p}}, \hat{\vec{x}}, X(t))$. We defined the Schrödinger picture by

$$\begin{aligned} \hat{\mathcal{H}}_{eff}(t) &\equiv U(t)^\dagger \hat{H}_{eff}(t) U(t) \\ &= \sum_n b_n^\dagger(0) \mathcal{E}_n(X(t)) b_n(0) - \sum_{n,m} b_n^\dagger(0) \langle n | i\hbar \frac{\partial}{\partial t} | m \rangle b_m(0) \end{aligned}$$

by introducing $U(t)$, $i\hbar \frac{\partial}{\partial t} U(t) = -\hat{H}_{eff}(t) U(t)$, with $U(0) = 1$.

lowest two levels, and we consider the familiar idealized model with only the lowest two levels. This simplification is expected to be valid for the analysis of the issues we are interested in. The effective Hamiltonian to be analyzed in the path integral (4) is then defined by the 2×2 matrix $h(X(t)) = (E_{nm}(X(t)))$. If one assumes that the level crossing takes place at the origin of the parameter space $X(t) = 0$, one needs to analyze the matrix

$$h(X(t)) = (E_{nm}(0)) + \left(\frac{\partial}{\partial X_k} E_{nm}(0) \right) X_k(t) \quad (11)$$

for sufficiently small $(X_1(1), X_2(1), \dots)$. By a time independent unitary transformation, which does not induce a geometric term, the first term is diagonalized. In the present approximation, essentially the four dimensional sub-space of the parameter space is relevant, and after a suitable re-definition of the parameters by taking linear combinations of $X_k(t)$, we write the matrix as [1]

$$h(X(t)) = \begin{pmatrix} E(0) + y_0(t) & 0 \\ 0 & E(0) + y_0(t) \end{pmatrix} + g\sigma^l y_l(t) \quad (12)$$

where σ^l stands for the Pauli matrices, and g is a suitable (positive) coupling constant.

The above matrix is diagonalized in a standard manner

$$h(X(t))v_{\pm}(y) = (E(0) + y_0(t) \pm gr)v_{\pm}(y) \quad (13)$$

where $r = \sqrt{y_1^2 + y_2^2 + y_3^2}$ and

$$v_+(y) = \begin{pmatrix} \cos \frac{\theta}{2} e^{-i\varphi} \\ \sin \frac{\theta}{2} \end{pmatrix}, \quad v_-(y) = \begin{pmatrix} \sin \frac{\theta}{2} e^{-i\varphi} \\ -\cos \frac{\theta}{2} \end{pmatrix} \quad (14)$$

by using the polar coordinates, $y_1 = r \sin \theta \cos \varphi$, $y_2 = r \sin \theta \sin \varphi$, $y_3 = r \cos \theta$. Note that $v_{\pm}(y(0)) = v_{\pm}(y(T))$ if $y(0) = y(T)$ except for $(y_1, y_2, y_3) = (0, 0, 0)$, and $\theta = 0$ or π . If one defines

$$v_m^\dagger(y) i \frac{\partial}{\partial t} v_n(y) = A_{mn}^k(y) \dot{y}_k \quad (15)$$

where m and n run over $\pm$, we have

$$\begin{aligned} A_{++}^k(y) \dot{y}_k &= \frac{(1 + \cos \theta)}{2} \dot{\varphi}, \\ A_{+-}^k(y) \dot{y}_k &= \frac{\sin \theta}{2} \dot{\varphi} + \frac{i}{2} \dot{\theta} = (A_{-+}^k(y) \dot{y}_k)^*, \\ A_{--}^k(y) \dot{y}_k &= \frac{(1 - \cos \theta)}{2} \dot{\varphi}. \end{aligned} \quad (16)$$

The effective Hamiltonian (10) is then given by

$$\begin{aligned} \hat{H}_{eff}(t) &= (E(0) + y_0(t) + gr(t)) b_+^\dagger b_+ \\ &+ (E(0) + y_0(t) - gr(t)) b_-^\dagger b_- - \hbar \sum_{m,n} b_m^\dagger A_{mn}^k(y) \dot{y}_k b_n. \end{aligned} \quad (17)$$

In the conventional adiabatic approximation, one approximates the effective Hamiltonian (17) by

$$\begin{aligned}\hat{H}_{eff}(t) \simeq & (E(0) + y_0(t) + gr(t))b_+^\dagger b_+ \\ & + (E(0) + y_0(t) - gr(t))b_-^\dagger b_- \\ & - \hbar[b_+^\dagger A_{++}^k(y)\dot{y}_k b_+ + b_-^\dagger A_{--}^k(y)\dot{y}_k b_-]\end{aligned}\quad (18)$$

which is valid for $Tgr(t) \gg \hbar\pi$, the magnitude of the geometric term. The Hamiltonian for b_- , for example, is then eliminated by a “gauge transformation”

$$b_-(t) = \exp\left\{-\frac{i}{\hbar} \int_0^t dt [E(0) + y_0(t) - gr(t) - \hbar A_{--}^k(y)\dot{y}_k]\right\} \tilde{b}_-(t) \quad (19)$$

in the path integral (8), and the amplitude $\langle 0|\hat{\psi}(T)b_-^\dagger(0)|0\rangle$, which corresponds to the probability amplitude in the first quantization, is given by (up to a wave function $\phi_E(\vec{x})$)

$$\begin{aligned}& \exp\left\{-\frac{i}{\hbar} \int_0^T dt [E(0) + y_0(t) - gr(t) - \hbar A_{--}^k(y)\dot{y}_k]\right\} \\ & \times v_-(y(T)) \langle 0|\tilde{b}_-(T)\tilde{b}_-^\dagger(0)|0\rangle\end{aligned}\quad (20)$$

with $\langle 0|\tilde{b}_-(T)\tilde{b}_-^\dagger(0)|0\rangle = 1$. For a 2π rotation in φ with fixed θ , for example, the geometric term gives rise to the well-known factor $\exp\{i\pi(1 - \cos\theta)\}$ by using (16) [1]. The corrections to the phase due to the finite $1/T$ may be evaluated by an iterative procedure [8], for example.

Another representation, which is useful to analyze the behavior near the level crossing point, is obtained by a further unitary transformation $b_m = U(\theta(t))_{mn}c_n$ where m, n run over $\pm$ with

$$U(\theta(t)) = \begin{pmatrix} \cos \frac{\theta}{2} & -\sin \frac{\theta}{2} \\ \sin \frac{\theta}{2} & \cos \frac{\theta}{2} \end{pmatrix}, \quad (21)$$

and the above effective Hamiltonian (17) is written as

$$\begin{aligned}\hat{H}_{eff}(t) &= (E(0) + y_0(t) + gr \cos \theta)c_+^\dagger c_+ \\ &+ (E(0) + y_0(t) - gr \cos \theta)c_-^\dagger c_- \\ &- gr \sin \theta c_+^\dagger c_- - gr \sin \theta c_-^\dagger c_+ - \hbar \dot{\varphi} c_+^\dagger c_+.\end{aligned}\quad (22)$$

In the above unitary transformation, an extra geometric term $-U(\theta)^\dagger i\hbar \partial_t U(\theta)$ is induced by the kinetic term of the path integral representation (8). One can confirm that this extra term precisely cancels the term containing $\dot{\theta}$ in $b_m^\dagger A_{mn}^k(y)\dot{y}_k b_n$ as in (16). We thus diagonalize the geometric terms in this representation. We also note that $U(\theta(T)) = U(\theta(0))$ if $X(T) = X(0)$ except for the origin, and thus the initial and final states receive the same transformation in scattering amplitudes.

In the infinitesimal neighborhood of the level crossing point, namely, for sufficiently close to the origin of the parameter space $(y_1(t), y_2(t), y_3(t))$ but $(y_1(t), y_2(t), y_3(t)) \neq (0, 0, 0)$, one may approximate (22) by

$$\begin{aligned}\hat{H}_{eff}(t) \simeq & (E(0) + y_0(t) + gr \cos \theta)c_+^\dagger c_+ \\ & + (E(0) + y_0(t) - gr \cos \theta)c_-^\dagger c_- - \hbar \dot{\varphi} c_+^\dagger c_+.\end{aligned}\quad (23)$$

To be precise, for any given *fixed* time interval T , $T\hbar\dot{\varphi} \sim 2\pi\hbar$ which is invariant under the uniform scale transformation $y_k(t) \rightarrow \epsilon y_k(t)$. On the other hand, one has $Tgr \sin \theta \rightarrow T\epsilon gr \sin \theta$ by the above scaling, and thus one can choose $T\epsilon gr \ll \hbar$. The terms $\pm gr \cos \theta$ in (23) may also be ignored in the present approximation.

In this new basis (23), the geometric phase appears only for the mode c_+ which gives rise to a phase factor $\exp\{i \int_C \dot{\varphi} dt\} = \exp\{2i\pi\} = 1$, and thus no physical effects. In the infinitesimal neighborhood of level crossing, the states spanned by (b_+, b_-) are transformed to a linear combination of the states spanned by (c_+, c_-) , which give no non-trivial geometric phases. The geometric terms are topological in the sense that they are invariant under the uniform scaling of $y_k(t)$, but their physical implications in conjunction with other terms in the effective Hamiltonian are not. For example, starting with the state $b_-^\dagger(0)|0\rangle$ one may first make $r \rightarrow \text{small}$ with fixed θ and φ , then make a 2π rotation in φ in the bases $c_\pm^\dagger|0\rangle$, and then come back to the original r with fixed θ and φ for a given fixed T ; in this cycle, one does not pick up any non-trivial geometric phase even though one covers the solid angle $2\pi(1 - \cos \theta)$. The transformation from $b_\pm$ to $c_\pm$ is highly non-perturbative, since a complete rearrangement of two levels is involved.

It is noted that one cannot simultaneously diagonalize the conventional energy eigenvalues and the induced geometric terms in (17) which is exact in the present two-level model (12). The topological considerations [3, 1] are thus inevitably approximate. In this respect, it may be instructive to consider a model without level crossing which is defined by setting

$$y_3 = \Delta E/2g \quad (24)$$

in (17), where ΔE stands for the minimum of the level spacing. The geometric terms then loose invariance under the uniform scaling of y_1 and y_2 . In the limit

$$\sqrt{y_1^2 + y_2^2} \gg \Delta E/2g, \quad (25)$$

(and thus $\theta \rightarrow \pi/2$), the geometric terms in (17) exhibit approximately topological behavior for the reduced variables (y_1, y_2) . Near the point where the level spacing becomes minimum, which is specified by

$$(y_1, y_2) \rightarrow (0, 0) \quad (26)$$

(and thus $\theta \rightarrow 0$), the geometric terms in (17) assume the same form of the geometric term as in (23). Our analysis above shows that the model with level crossing exhibits precisely the same topological properties for any finite T .

It is instructive to analyze an explicit example in Refs. [13, 14] where the following parametrization has been introduced

$$(y_1, y_2, y_3) = (B_0(b_1 + \cos \omega t), B_0 \sin \omega t, B_z) \quad (27)$$

and $g = \mu$. The case $b_1 = 0$ and $B_z \neq 0$ corresponds to the model without level crossing discussed above, and the geometric phase becomes trivial for $B_0 \rightarrow 0$. The case $b_1 = B_z = 0$ describes the situation in (23), namely, a closed cycle in the infinitesimal neighborhood

of level crossing for $B_0 \rightarrow 0$ with $T = 2\pi/\omega$ kept fixed, and the geometric phase becomes trivial. On the other hand, the usual adiabatic approximation (18) (with $\theta = \pi/2$ in the present model) in the neighborhood of level crossing is described by $b_1 = B_z = 0$ and $B_0 \rightarrow 0$ (and $\omega \rightarrow 0$) with

$$\mu B_0/\hbar\omega \gg 1 \quad (28)$$

kept fixed, namely, the effective magnetic field is always strong; the topological proof of phase-change rule [3] is based on the consideration of this case. It should be noted that the geometric phase becomes trivial for $B_0 \rightarrow 0$ with $b_1 = B_z = 0$ and

$$\mu B_0/\hbar\omega \ll 1 \quad (29)$$

kept fixed. (If one starts with $b_1 = B_z = 0$ and $\omega = 0$, of course, no geometric terms.) It is clear that the topology is non-trivial only for a quite narrow (essentially measure zero) window of the parameter space (B_0, ω) in the approach to the level crossing $B_0 \rightarrow 0$. In this analysis, it is important to distinguish the level crossing problem from the motion of a spin 1/2 particle; the wave functions (14) are single valued for a 2π rotation in φ with fixed θ .

The path integral (4), where the Hamiltonian is diagonalized both at $t = 0$ and $t = T$ if $X(T) = X(0)$, shows no obvious singular behavior at the level crossing point. On the other hand, the path integral (8) is subtle at the level crossing point; the bases $\{v_n(\vec{x}, X(t))\}$ are singular on top of level crossing as in (14), and thus the unitary transformation U to (8) and the induced geometric terms become singular there. The present analysis suggests that the path integral is not singular for any finite T , as is expected from (4). We consider that this result is natural since the starting Hamiltonian (1) does not contain any obvious singularity.

The conventional treatment of geometric phases in the (precise) adiabatic approximation is based on the premise that one can choose T sufficiently large for any *given* $\epsilon \sim r$ such that

$$Tg\epsilon \gg \hbar, \quad (30)$$

and thus $T \rightarrow \infty$ for $\epsilon \rightarrow 0$, namely, it takes an infinite amount of time to approach the level crossing point [1, 2]. Finite T may however be appropriate in practical applications, as is noted in [1]. Because of the uncertainty principle $T\Delta E \geq \frac{1}{2}\hbar$, the (physically measured) energy uncertainty for any given fixed T is not much different from the magnitude of the geometric term $2\pi\hbar$, and the level spacing becomes much smaller than these values in the infinitesimal neighborhood of level crossing for the given T . An intuitive picture behind (23) is that the motion in φ smears the “monopole” singularity for arbitrarily large but finite T .

The notion of Berry’s phase is known to be useful in various physical contexts [15]-[16], and the topological considerations are often crucial to obtain a qualitative understanding of what is going on. Our analysis however shows that the geometric phase associated with level crossing becomes topologically trivial in practical physical settings with any finite T . This is in sharp contrast to the Aharonov-Bohm phase [7] which is induced by the

time-independent gauge potential and topologically exact for any finite time interval T . The similarity and difference between the geometric phase associated with level crossing and the Aharonov-Bohm phase have been recognized in the early literature [1, 7]. Also, the correction to the geometric phase in terms of the small slowness parameter ϵ has been analyzed, and the closer to a degeneracy a system passes the slower is the necessary passage for adiabaticity has been noted in [8]. But, to our knowledge, the fact that the geometric phase becomes topologically trivial for practical physical settings with any fixed finite T , such as in the practical Born-Oppenheimer approximation where a large but finite ratio of two time scales is involved, has not been clearly stated in the literature. We emphasize that this fact is proved independently of the adiabatic approximation.

The notion of the geometric phase is useful, but great care needs to be exercised as to its topological properties.

References

- [1] M.V. Berry, Proc. Roy. Soc. **A392**, 45 (1984).
- [2] B. Simon, Phys. Rev. Lett. **51**, 2167 (1983).
- [3] A.J. Stone, Proc. Roy. Soc. **A351**, 141 (1976).
- [4] H. Longuet-Higgins, Proc. Roy. Soc. **A344**, 147 (1975).
- [5] F. Wilczek and A. Zee, Phys. Rev. Lett. **52**, 2111 (1984).
- [6] H. Kuratsuji and S. Iida, Prog. Theor. Phys. **74**, 439 (1985).
- [7] Y. Aharonov and J. Anandan, Phys. Rev. Lett. **58**, 1593 (1987).
- [8] M.V. Berry, Proc. Roy. Soc. **A414**, 31 (1987).
- [9] N. Manini and F. Pistolesi, Phys. Rev. Lett. **85**, 3067 (2000).
- [10] N. Mukunda, Arvind, E. Ercolessi, G. Marmo, G. Morandi, and R. Simon, Phys. Rev. **A67**, 042114 (2003).
- [11] Y. Hasegawa, R. Loidl, M. Baron, G. Badurek, and H. Rauch, Phys. Rev. Lett. **87**, 070401 (2001).
- [12] Y. Hasegawa, R. Loidl, M. Baron, N. Manini, F. Pistolesi, and H. Rauch, Phys. Rev. **A65**, 052111 (2002).
- [13] Y. Lyanda Geller, Phys. Rev. Lett. **71**, 657 (1993).
- [14] R. Bhandari, Phys. Rev. Lett. **88**, 100430 (2002).
- [15] A. Shapere and F. Wilczek, ed., *Geometric Phases in Physics* (World Scientific, Singapore, 1989), and papers reprinted therein.

- [16] As for a recent comprehensive account of this subject see, for example, D. Chruscinski and A. Jamiolkowski, *Geometric Phases in Classical and Quantum Mechanics* (Birkhauser, Berlin, 2004).